

At two months old:

- Diphtheria, tetanus, pertussis (whooping cough), polio and Haemophilus influenzae type b (Hib) (DTaP/IPV/Hib) - **one injection**
- Pneumococcal infection - pneumococcal conjugate vaccine (PCV) - **one injection**

At three months old:

- Diphtheria, tetanus, pertussis (whooping cough), polio and Haemophilus influenzae type b (Hib) (DTaP/IPV/Hib) - **one injection**
- Meningitis C (meningococcal group C) (MenC) - **one injection**

At four months old:

- Diphtheria, tetanus, pertussis (whooping cough), polio and Haemophilus influenzae type b (Hib) (DTaP/IPV/Hib) - **one injection**
- Meningitis C (meningococcal group C) (MenC) - **one injection**
- Pneumococcal infection - pneumococcal conjugate vaccine (PCV) - **one injection**

At around 12 months old:

- Haemophilus influenzae type b (Hib) and meningitis C (Hib/MenC) - booster dose in **one injection**

At around 13 months old:

- Measles, mumps and rubella (German measles) (MMR) - **one injection**
- Pneumococcal infection - pneumococcal conjugate vaccine (PCV) - **one injection**

Three years four months to five years old (pre-school):

- Diphtheria, tetanus, pertussis (whooping cough) and polio (dTaP/IPV or DTaP/IPV) - **one injection**
- Measles, mumps and rubella (German measles) (MMR) - **one injection**

13 to 18 years old:

- Diphtheria, tetanus and polio (Td/IPV) - **one injection**